



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Review

Role of poly(ADP-ribose) polymerases in the regulation of inflammatory processes

Péter Bai^{*}, László Virág^{*}

Department of Medical Chemistry, Medical and Health Science Center, University of Debrecen, Debrecen H-4032, Hungary
 Cell Biology and Signaling Research Group of the Hungarian Academy of Sciences, Debrecen, Hungary

ARTICLE INFO

Article history:

Received 8 August 2012

Revised 6 September 2012

Accepted 16 September 2012

Available online 26 September 2012

Edited by Laszlo Nagy

Keywords:

Inflammation

Allergy

Oxidative stress

Nitrosative stress

Chemokine

Adhesion molecule

Gene expression

ABSTRACT

PARP enzymes influence the immune system at several key points and thus modulate inflammatory diseases. PARP enzymes affect immune cell maturation and differentiation and regulate the expression of inflammatory mediators such as cytokines, chemokines, inducible nitric oxide synthase and adhesion molecules. Moreover, PARP enzymes are key regulators of cell death during inflammation-related oxidative and nitrosative stress. Here we provide an overview of the different inflammatory diseases regulated by PARP enzymes.

© 2012 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

1. Introduction

Poly(ADP-ribose) polymerases (PARPs) constitute a family of 17 enzymes [1] sharing a highly conserved PARP signature motif in the catalytic domain. The prototypical PARP enzymes (PARP-1 and -2) possess a DNA binding domain enabling them to bind to damaged DNA and thus become activated [2]. Active PARP-1 and PARP-2 synthesize poly(ADP-ribose) polymers from NAD⁺ marking proteins near the DNA damage site. Although the characterization of the newly discovered PARP enzymes has not yet been accomplished, it is already apparent that PARP enzymes perform a wide variety of tasks ranging from DNA repair, regulation of gene expression to cell death [3].

The anti-inflammatory properties of PARP inhibitors have been recognized over a decade [4] and the number of publications on these effects is still on the rise. Our aim in the present chapter is to provide an overview of the complex role of PARP enzymes in inflammatory processes ranging from immune cell maturation to inflammatory gene expression and to explain the mechanism of protection provided by the PARP knockout phenotypes or by PARP inhibition in inflammatory tissue injury.

2. Poly(ADP-ribose) polymerases in the maturation and differentiation of immune cells

In adult mammals immune cells are formed in the bone marrow by the division of resident stem cells. These premature immune cells then travel to different immune organs (thymus, the Peyer's patches of the gastrointestinal (GI) tract, etc.) and undergo a process called maturation in order to yield competent, functional, non-self reactive cells that can then exit to the periphery and perform adequate immune functions. Although no systematic investigation took place as of yet, current data suggest that different PARP enzymes are necessary for the maturation and differentiation of different immune cell types.

An important point during T and B cell development is the VDJ recombination that is responsible for the generation of a wide repertoire of immunoglobulins and T cell receptors. VDJ recombination involves DNA rearrangement and thus the resolution and rejoining of DNA strand breaks in which the elementary role of the DNA repair machinery has been shown [5] prompting research to assess the role of PARPs in these processes. Only minor alterations have been reported in B cells of PARP-1 knockout mice. However, PARP inhibition increased the frequency of Ig class switch [6] leading to variations in immunoglobulin class composition [7]. Moreover, the deficient VDJ recombination in the T cells of SCID (severe combined immunodeficiency) mice could be rescued by knocking out PARP-1 [8] indicating the antirecombinogenic effect of PARP-1 during T cell maturation.

^{*} Corresponding authors at: Department of Medical Chemistry, University of Debrecen, Nagyterdei krt 98, Pf. 7, Debrecen H-4032, Hungary. Fax: +36 52 412 566.
 E-mail addresses: baip@med.unideb.hu (P. Bai), lvirag@med.unideb.hu (L. Virág).

PARP-1 influences mature B cell differentiation and proliferation by binding to the *Bcl6* gene in a sequence-specific manner acting as a transcriptional repressor. Since *Bcl6* expression is restricted to germinal centers, where active cell division takes place, PARP-1 binding and activation is likely to be necessary for B cells to exit from germinal centers [9].

Both T cell function and development may be affected by PARP-1 and -2. A recent report has described an increased number but normal phenotype and function of *Foxp3*⁺ regulatory T cells and consequently impaired proliferation of CD4⁺ T cells and suppressed IL-2 production in PARP-1^{-/-} mice [10]. This may partly explain suppressed inflammation observed in PARP-1 knockout mice in models of T cell-mediated inflammatory diseases. T cell-dependent antibody responses were also found to be impaired in PARP-1^{-/-} mice [7]. PARP-2 also affects T cell number through impairing thymopoiesis. PARP-2 was abundantly expressed in the subcapsular zone of the thymus with decreased expression towards the center of the thymus suggestive of its involvement in thymocyte proliferation [11]. Indeed, PARP-2 deficient thymocytes die via p53-mediated apoptosis that leads to decreased number of CD4⁺, CD8⁺ thymocytes in the thymus [12,13].

Impaired T cell-mediated inflammation can thus be attributed to several factors, such as impaired inflammatory gene expression (discussed in the next chapter) and difference in the composition of T cell populations. Scattered data in the literature indicate that the differentiation of other inflammatory cell types, such as monocytes and dendritic cells may also be affected by PARP-1 [14,15].

3. Molecular events in inflammation regulated by poly(ADP-ribose) polymerases

3.1. Transcription factors

PARP enzymes interact with a large number of transcription factors of which many are involved in the regulation of inflammatory gene expression. The first one identified was NFκB [16] providing an explanation at the molecular level for the protection of PARP-1 knockout mice from endotoxin shock. Several other transcription factors and cofactors involved in inflammatory regulation such as NFAT [17,18], AP-1 [19–22], YY1 [23], Sp1 [24], SIRT1 [25,26] were also found to be modulated by PARP-1. Recent data pointed towards the involvement of other PARP enzymes in the transcriptional regulation of inflammatory processes. PARP-2 regulates the activity of several members of the peroxisome proliferator activated receptor family (PPARs) [27], or SIRT1 [28,29], while PARP-14 mediates STAT6 transcription [30,31]. However, the relevance of these latter findings in inflammations requires further investigation.

It is interesting to note that most transcription factors listed above are also activated by oxidative stress, thus considered redox-sensitive. It is debated whether PARP activity is necessary for the proper activation of these transcription factors since *in vitro* data suggest that DNA binding and PARP activity might be dispensable for NFκB activation [32]. In contrast, a large set of *in vivo* data support the anti-inflammatory effects of PARP inhibitors. Recently two excellent studies linked PARP1 to upstream events of NFκB pathway by demonstrating that PARP-1 activation is necessary for the ubiquitination and degradation of the inhibitor of NFκB kinase (IKK) promoting NFκB activation [33,34]. The actual mode of how PARP-1 contributes to the activation of NFκB and other transcription factors in different models will require further investigation.

3.2. Cytokines and chemokines

Inflammatory cytokines such as TNFα, IL-1β, IL-6 and IL-12 are central regulators of the inflammatory process orchestrated by Th1 cells. These cytokines boost the expression of other cytokines and

chemokines, adhesion molecules, matrix metalloproteinases, inducible nitric oxide synthase (iNOS), and cyclooxygenase-2 (COX-2). PARP inhibition or the PARP-1 knockout genotype has been shown to result in suppressed levels of the inflammatory cytokines in various animal models of inflammation. Similarly, many chemoattractant chemokines (e.g. IL-8, MIP-1, MIP-2, MCP-1) were also found to be downregulated in animals lacking a functional PARP-1 gene or treated with a PARP inhibitor. These changes may represent the underlying mechanism for the inhibition of inflammatory cell migration as reported in many Th1-mediated inflammatory conditions.

PARPs other than PARP1 may also alter the production of chemokines and cytokines. Deletion of PARP-2 leads to alterations in the transcriptome, notably in the expression of genes involved in inflammatory regulation [reviewed in [35]]. Interestingly there seems to be an overlap with the inflammatory genes dysregulated upon PARP-1 ablation: e.g. IL-1β and TNFα levels were lowered both in the absence of PARP-1 or PARP-2 [36,37]. Moreover, knocking out PARP-14 enhances IL-4-induced STAT6-mediated gene expression that impacts on immune processes notably in Th2-mediated processes in the lungs [30,31].

3.3. Adhesion factors

Another important group of regulated proteins include different adhesion proteins, such as cell adhesion molecules (I-CAM, V-CAM, L-CAM) and selectins. Suppressed expression of adhesion molecules on the surface of endothelial and immune cells decreases the migration of inflammatory cells to the site of inflammation and thus inhibits inflammation. In the absence, or upon inhibition of PARP-1, the expression of these adhesion molecules is depressed [21,38]. This effect may have a dual cause: on one hand NFκB directly regulates the expression of these genes, while on the other hand, a secondary loop may exist, where the NFκB-regulated cytokines regulate the expression of the adhesion molecules [39,40].

3.4. Inflammatory mediators: iNOS, COX-2 and MMPs

An important inflammatory gene regulated by PARP-1 is iNOS that is responsible for NO synthesis under inflammatory conditions [41,42]. NO derived from iNOS may combine with superoxide to form peroxynitrite [43–45]. Interestingly, iNOS, under certain pathological conditions, where iNOS substrates are limiting, may become uncoupled and produce superoxide instead of nitrogen monoxide [46]. This mechanism has been suggested to enhance peroxynitrite production and tissue damage [47]. Although it is a rather labile reactive nitrogen species, peroxynitrite can travel through biological compartments, can diffuse through membranes and can reach the nucleus to trigger oxidative and nitrative damage to DNA, proteins and lipids [48].

As iNOS is regulated via NFκB activation, therefore PARP inhibition is capable of markedly reducing iNOS expression and nitrosative stress [21,39]. Cyclooxygenase-2 (COX-2) is also involved in generating free radicals and has been shown to be regulated by NFκB. In the case of COX-2, however, a more direct regulatory link may be more important: PARP-1 has been demonstrated to bind to the mouse COX-2 promoter region through interactions with the inhibitory element resulting in the inhibition of COX-2 expression [49].

Increased oxidative stress is capable of inducing redox-sensitive enzymes, such as matrix metalloproteinases (MMPs). In addition to proteolytic cleavage, MMPs can also be activated by free radical-induced structural changes [50] and are indispensable for cellular movements within tissues. Several studies have shown that MMP activation during inflammation (e.g. MMP-9 activation in skin inflammation) can be prevented by knocking out PARP-1 or by




Fig. 1. PARP as a central regulator of inflammatory pathways. DNA damage triggers the activation of PARP-1 and -2. PARP activation has proinflammatory properties through lowering the number of inhibitory T cells, enhancing antigen presentation by dendritic cells. Moreover, oxidative stress stimulates redox-sensitive transcription factors that are co-activated by PARP-1. These transcription factors regulate the expression of chemokines, adhesion molecules, iNOS, MMP and COX-2. These effects lead to inflammatory cell homing, tissue invasion and activation which in turn further enhances inflammation-related oxidative stress and a vicious, self-intensifying cycle of inflammation.

pharmacological PARP inhibition [21,51–54]. Moreover, tissue inhibitors of MMPs (TIMPs) that keep MMP activity under control are inversely regulated with MMPs [21,51,54]. It has not been determined as of yet whether these effects are related to modulation of MMP and TIMP gene expression by PARylation or PARylation controls MMP activation via regulation of cellular redox status.

From the above it is clear that PARP-1 plays a central role in inflammation by influencing multiple key events such as the production of cytokines, expression of adhesion molecules, iNOS and COX-2 and probably MMPs and TIMPs through modulating different transcription factors (Fig. 1.). These events form a complex signaling network in which the proteins display a multilevel interrelationship with each other. For example NF κ B enhances iNOS expression, hence oxidative stress that in turn activates both NF κ B and PARP-1/2, creating a vicious, self intensifying cycle. Moreover, the PARP-1/2- and NF κ B-dependent expression of adhesion factors and MMPs contribute to inflammatory cell migration and thus to oxidative stress which feeds back to both PARP-1/2 and NF κ B. This high level of complexity makes the *in vivo* dissection of causal relationships challenging. In the same time this central role of PARP-1 in inflammatory circuitry makes these enzymes promising targets of anti-inflammatory drug development.

4. Oxidative stress-induced activation of poly(ADP-ribose) polymerases in inflammation

Inflammatory processes are associated with high level of oxidative stress. The oxidants produced for example by iNOS, NADPH oxidases, COX-2 or the mitochondrial respiratory chain cause excessive DNA damage. Berger and colleagues [55] proposed that severe DNA damage may lead to excessive PARP activation which may deplete cellular NAD⁺ pools compromising cellular metabolism. Under such conditions glycolytic flux slows down [56] thereby impairing an oxygen-insensitive metabolic pathway. Moreover,

NAD⁺ depletion turns on NAD⁺ resynthesis via nicotinamide mononucleotide adenylyltransferase (NMNAT) and phosphoribosyl pyrophosphate synthetase (PPS) at the expense of ATP. The often marked NAD⁺ depletion may lead to consumption of cellular ATP [3,55]. Under low ATP, low NAD⁺ conditions, mitochondrial F1/F0 ATPase turns to function as ATPase rather than ATP synthase [57] further deteriorating energy depletion. Compromised cellular energetics may contribute to the pathological sequelae of excessive poly(ADP-ribosyl)ation in inflammations.

In severe oxidative stress, cellular energy status also affects cell death pathways. Failure of DNA repair initiates apoptotic cell death. Since apoptosis also relies on ATP, cleavage of PARP-1 and -2 by caspase-3 and -7 may be viewed as a preventive measure aiming at the conservation of cellular ATP [58,59]. During apoptosis cellular content is crosslinked and cells disintegrate into membrane-coated vesicles marked with the “eat-me” signal phosphatidyl-serine for phagocytosis. These vesicles are then cleared by infiltrating macrophages and resident, neighboring cells, therefore no leakage of cellular content occurs under apoptosis.

If the apoptotic process cannot be executed (e.g. due to lack of sufficient amount of ATP or inhibition of caspases), then cells may die by necrosis. Necrosis is characterized by the lack of the morphological and biochemical hallmarks of apoptosis. Most notably, plasma membrane integrity is compromised early during necrosis and cellular content may leak into the environment. Such leakage may cause secondary tissue injury by causing further inflammation. Initially necrosis was thought to be an uncontrolled form of cell death. However, recently programmed features and necrotic signaling molecules (RIP kinases, non-caspase proteases, cyclophylline D) have been identified in necrosis indicating that necrosis is not necessarily beyond regulatory control [60]. Overactivation of PARP-1 has been increasingly recognized as a mediator of necrosis that is highly amenable to pharmacological intervention. In severe oxidative stress for example PARP-1-mediated necrosis serves as a

backup mechanism eliminating severely injured cells even when the apoptotic machinery is incapacitated. In fact PARP activation actively switches the default apoptotic cell death towards necrosis. When PARP activity is inhibited or *parp-1* gene is knocked out, necrosis can be reverted to apoptosis [3,61–63].

5. Inflammatory diseases mediated by poly(ADP-ribose) polymerases

Previously we reviewed the molecular determinants for the proinflammatory role of PARylation. These mechanisms have been validated in a vast array of inflammatory conditions identifying a large number of diseases where PARP inhibition may provide therapeutic benefits (detailed in Table 1.). It is important to note that almost all organs display inflammatory conditions that can be attenuated by PARP inhibition indicating that PARP affects central events in inflammatory signaling. The diseases listed in Table 1 are quite diverse in terms of etiology, pathomechanism (infectious, autoimmune, allergic, irritation-induced, etc.) being organ-restricted versus generalized inflammations. Most of the conditions listed are mediated by IL-2- and IFN γ -producing Th1 cells. Others such as asthma are typical Th2-mediated conditions dominated by the production of Th2 cytokines such as IL-4, -5 and -14. Recent reinterpretation of Th1/Th2 dichotomy as triggered by the identification of the novel Th17 cell type shed light on the complexity of immunoregulation in inflammatory diseases such as collagen-induced arthritis (CIA) and experimental allergic encephalomyelitis (EAE), animal models of the human diseases rheumatoid arthritis and multiple sclerosis, respectively. These latter two models are now considered as Th17 rather than Th1-mediated conditions. (The new Th17 cell type is characterized by the production of IL17 and 23.)

Since PARP-1 regulates such central events of inflammatory signaling as the activation of NF κ B and AP-1 transcription factors,

therefore it is not surprising that PARP inhibition has beneficial effects in a wide spectrum of inflammatory conditions. Nonetheless, the mechanism of the beneficial effects of PARP inhibition may differ between Th1 and Th2-mediated conditions. Whereas the NF κ B co-activator function of PARP-1 has been made responsible for the effects of PARP inhibitors in Th1 diseases (and this may also contribute to the pathology of Th2-mediated conditions), recently a novel mechanism has been described for asthma, a prototypical Th2-mediated disease. Datta et al. [64] have demonstrated in a murine model of asthma that inhibition of eosinophil migration in PARP-1^{-/-} mice is controlled at a step downstream of IL-4 and upstream of IL-5. PARP-1 was found to stabilize STAT-6 protein (without affecting mRNA) in an allergen-stimulation dependent manner. STAT-6, a central regulator of IL-5 expression undergoes degradation in the absence of functional PARP-1. This example nicely illustrates that the role played by PARPs in inflammation is complex and may differ in various forms of inflammation.

An inflammatory component has recently been identified as an important pathogenetic factor in the development of several chronic diseases of the cardiovascular system such as atherosclerosis, chronic heart failure and cardiovascular aging [65]. Both PARP inhibition and knockout of PARP-1 have been shown to have beneficial effects in murine models of these conditions [66–70]. Although the molecular mechanisms underlying these positive effects are not well understood, it is plausible to hypothesize that the anti-inflammatory effects of PARP inhibition may contribute to improved clinical parameters observed in these models. PARP activation is associated with numerous human inflammatory diseases such as allograft rejection, various forms of shock, arthritis, colitis, diabetic complications, transplant rejection, COPD [3,71,72] opening the door for the possible clinical use of PARP inhibitors in these diseases. Several clinical trials have been conducted with PARP inhibitors demonstrating their good tolerability [73,74] and their potential as anti-cancer agents [75]. Moreover, in the study

Table 1
Inflammatory diseases mediated by poly(ADP-ribosyl)ation. PARPi – PARP inhibition, KO – knockout, all other abbreviations in text.

Organ	Disease	Protection provided by	Ref.
Multiorgan diseases	Sepsis	PARPi	[82]
	Endotoxic shock	PARP-1 K.O., PARPi	[16,83]
	Heamorrhagic shock	PARP-1 K.O.	[84]
Central nervous system	Allergic encephalomyelitis	PARPi	[85]
	Meningitis-related central nervous system complications	PARP-1 K.O., PARPi	[86]
	Myelitis	PARPi	[87]
	Astrocyte activation	PARP-1 KO PARP-2 KO PARP-3 KO	[36]
Joints	Arthritis		[88]
Bones	Temporomandibular joint disorder	PARPi	[89]
	Cherubism	Tankyrase activation	[79]
GI tract	Colitis	PARP-1 and -2 K.O.	[37,38]
	Pancreatitis	PARP-1 K.O. (PARP-2 has no effect), PARPi	[90]
	Peritonitis	PARP-1 K.O.	[91]
	Asthma	PARPi	[64,92–94]
Respiratory system		PARP-14 activation	[31]
	COPD	–	[95]
	Pleuritis	PARPi	[96]
	ARDS	PARP-1 K.O., PARPi	[97]
	Ventilation-induced inflammation	PARPi	[98]
	Rejection of tracheal allografts	PARPi	[99]
	Contact hypersensitivity	PARP-1 K.O., PARPi	[21,51,100]
Skin	Irritative dermatitis	PARP-1 K.O., PARPi	[21,51]
	Sunburn-related dermal inflammation	PARPi	[101]
	Uveitis	PARPi	[102]
Eye			
Ear	Cochlear lateral wall damage	PARP-1 K.O.	[103]
Skeletal muscle	Movement-induced damage associated inflammation		[104]
Kidneys	Renal hypertrophy	PARPi	[105]
	Cisplatin-induced kidney inflammation	PARPi, PARP-1 K.O.	[106]
	Diabetic nephropathy	PARPi	[107,108]
Cardiovascular system	Atherosclerosis	PARPi	[54,109–111]

utilizing the PARP inhibitor INO1001, Morrow and colleagues have shown that administration of INO1001 resulted in a tendency to reduce IL-6 and CRP levels pointing towards suppressed inflammation [73].

In addition to the beneficial effects of PARP inhibitors and protection provided by knockout phenotypes, the role of PARPs in inflammation is also indicated by association of certain PARP-1 promoter haplotypes and SNPs with diseases. Such associations have been reported for systemic lupus erythematosus [76], autoimmune nephritis [76] or rheumatoid arthritis [77] further supporting a possible link between PARP-1 and autoimmune inflammation.

6. Conclusion

As detailed above, PARP activation is a key event in the propagation of inflammation and PARP inhibition suppresses inflammation at multiple molecular events (for overview, see Fig. 1). Key components of the anti-inflammatory effects of PARP inhibition are the following: (1) Members of the PARP family may be necessary for the maturation and differentiation of cells of the immune system with PARP-1 and -2 playing a role in the regulation of T cell function. (2) PARPs are integral and central players of the self intensifying circle of inflammation by regulating the expression of inflammatory mediators such as cytokines, chemokines, adhesion molecules, iNOS and COX-2. A key molecular event in this regulation is the interaction of PARP-1 with transcription factors such as NF κ B and AP-1. (3) Free radicals may cause DNA damage and activate PARP-1 and -2. In case of severe DNA damage, PARP over-activation may cause cell dysfunction and cell death which may contribute to inflammation accompanying severe conditions such as shock and ischemia–reperfusion injuries.

Several open questions concerning PARP-related inflammatory diseases remain to be resolved. These include but are not limited to the mechanism of gender bias described in the role of PARP in inflammations [78]. Furthermore, it will be important to determine the involvement of further members of the PARP family in the regulation of inflammatory processes. Some of them have already been shown to associate with inflammatory pathologies: such as PARP-2 (reviewed in [35]), PARP-3 [36], tankyrases [79], or PARP-14 [30,31]. Moreover, poly(ADP-ribose) glycohydrolase has been implicated in the regulation of immune functions [80,81] underlining the central role of poly(ADP-ribosyl)ation in inflammation. A key question is whether the powerful anti-inflammatory effects of PARP inhibitors as observed in animal models can be translated to human diseases and to determine the risk benefit ratio of acute and chronic administration of PARP inhibitors in non-cancer patients.

Acknowledgements

This work was supported by Bolyai fellowship to P.B., Grants from the National Innovation Office (TÉT_09-2010-0023, Baross program Seahorse Grant, TÁMOP-4.2.1/B-09/KONV-2010-0007, TÁMOP-4.2.2. A-11/1/KONV-2012-0025, TÁMOP-4.2.2/B-10/1-2010-0024, TÁMOP-4.2.2/B-10/1-2010-0024), from OTKA (CNK80709, K82009, K75864, PD83473, K105872) and from the University of Debrecen, Medical and Health Science Center (Mec-8/2011).

References

- [1] Ame, J.C., Spelnhauer, C. and de Murcia, G. (2004) The PARP superfamily. *BioEssays* 26, 882–893.
- [2] Benjamin, R.C. and Gill, D.M. (1980) Poly(ADP-ribose) synthesis in vitro programmed by damaged DNA. A comparison of DNA molecules containing different types of strand breaks. *J. Biol. Chem.* 255, 10502–10508.
- [3] Virag, L. and Szabo, C. (2002) The therapeutic potential of poly(ADP-ribose) polymerase inhibitors. *Pharmacol. Rev.* 54, 375–429.
- [4] Weltin, D., Picard, V., Aupeix, K., Varin, M., Oth, D., Marchal, J., Dufour, P. and Bischoff, P. (1995) Immunosuppressive activities of 6(5H)-phenanthridinone, a new poly(ADP-ribose)polymerase inhibitor. *Int. J. Immunopharmacol.* 17, 265–271.
- [5] Taccioli, G.E., Rathbun, G., Oltz, E., Stamato, T., Jeggo, P.A. and Alt, F.W. (1993) Impairment of V(D)J recombination in double-strand break repair mutants. *Science* 260, 207–210.
- [6] Shockett, P. and Stavnezer, J. (1993) Inhibitors of poly(ADP-ribose) polymerase increase antibody class switching. *J. Immunol.* 151, 6962–6976.
- [7] Ambrose, H.E., Willmott, S., Beswick, R.W., Dantzer, F., de Murcia, J.M., Yelamos, J. and Wagner, S.D. (2009) Poly(ADP-ribose) polymerase-1 (Parp-1)-deficient mice demonstrate abnormal antibody responses. *Immunology* 127, 178–186.
- [8] Morrison, C., Smith, G.C., Stingl, L., Jackson, S.P., Wagner, E.F. and Wang, Z.Q. (1997) Genetic interaction between PARP and DNA-PK in V(D)J recombination and tumorigenesis. *Nat. Genet.* 17, 479–482.
- [9] Ambrose, H.E., Papadopoulos, V., Beswick, R.W. and Wagner, S.D. (2007) Poly(ADP-ribose) polymerase-1 (Parp-1) binds in a sequence-specific manner at the Bcl-6 locus and contributes to the regulation of Bcl-6 transcription. *Oncogene* 26, 6244–6252.
- [10] Nasta, F., Laudisi, F., Sambucci, M., Rosado, M.M. and Pioli, C. (2010) Increased Foxp3+ regulatory T cells in poly(ADP-Ribose) polymerase-1 deficiency. *J. Immunol.* 184, 3470–3477.
- [11] Schreiber, V., Ame, J.C., Dolle, P., Schultz, I., Rinaldi, B., Fraulob, V., Menissier-de Murcia, J. and de Murcia, G. (2002) Poly(ADP-ribose) polymerase-2 (PARP-2) is required for efficient base excision DNA repair in association with PARP-1 and XRCC1. *J. Biol. Chem.* 277, 23028–23036.
- [12] Yelamos, J. et al. (2006) PARP-2 deficiency affects the survival of CD4+CD8+ double-positive thymocytes. *EMBO J.* 25, 4350–4360.
- [13] Nicolas, L. et al. (2010) Loss of poly(ADP-ribose) polymerase-2 leads to rapid development of spontaneous T-cell lymphomas in p53-deficient mice. *Oncogene* 29, 2877–2883.
- [14] Selle, A., Ullrich, O., Harnacke, K. and Hass, R. (2007) Retrodifferentiation and rejuvenation of senescent monocytic cells requires PARP-1. *Exp. Gerontol.* 42, 554–562.
- [15] Aldinucci, A. et al. (2007) A key role for poly(ADP-ribose) polymerase-1 activity during human dendritic cell maturation. *J. Immunol.* 179, 305–312.
- [16] Oliver, F.J. et al. (1999) Resistance to endotoxic shock as a consequence of defective NF- κ B activation in poly(ADP-ribose) polymerase-1 deficient mice. *EMBO J.* 18, 4446–4454.
- [17] Olabisi, O.A. et al. (2008) Regulation of transcription factor NFAT by ADP-ribosylation. *Mol. Cell. Biol.* 28, 2860–2871.
- [18] Valdor, R. et al. (2008) Regulation of NFAT by poly(ADP-ribose) polymerase activity in T cells. *Mol. Immunol.* 45, 1863–1871.
- [19] Zingarelli, B., Hake, P.W., Burroughs, T.J., Piraino, G., O'Connor, M. and Denenberg, A. (2004) Activator protein-1 signalling pathway and apoptosis mediated by poly(ADP-ribose) polymerase-1 in experimental colitis. *Immunology* 113, 509–517.
- [20] Ha, H.C. (2004) Defective transcription factor activation for proinflammatory gene expression in poly(ADP-ribose) polymerase 1-deficient glia. *Proc. Natl. Acad. Sci. USA* 101, 5087–5092.
- [21] Brunyanszki, A. et al. (2010) Genetic ablation of PARP-1 protects against oxazolone-induced contact hypersensitivity by modulating oxidative stress. *J. Invest. Dermatol.* 130, 2629–2637.
- [22] Erdelyi, K. et al. (2009) Dual role of poly(ADP-ribose) glycohydrolase in the regulation of cell death in oxidatively stressed A549 cells. *FASEB J.* 23, 3553–3563.
- [23] Oei, S.L., Griesenbeck, J., Schweiger, M., Babich, V., Kropotov, A. and Tomilin, N. (1997) Interaction of the transcription factor YY1 with human poly(ADP-ribosyl) transferase. *Biochem. Biophys. Res. Commun.* 240, 108–111.
- [24] Ha, H.C., Hester, L.D. and Snyder, S.H. (2002) Poly(ADP-ribose) polymerase-1 dependence of stress-induced transcription factors and associated gene expression in glia. *Proc. Natl. Acad. Sci. USA* 99, 3270–3275.
- [25] Bai, P. et al. (2011) PARP-1 inhibition increases mitochondrial metabolism through SIRT1 activation. *Cell Metab.* 13, 461–468.
- [26] Bai, P. and Canto, C. (2012) The role of PARP-1 and PARP-2 enzymes in metabolic regulation and disease. *Cell Metab.* 16, 290–295.
- [27] Bai, P. et al. (2007) Poly(ADP-ribose) polymerase-2 controls adipocyte differentiation and adipose tissue function through the regulation of the activity of the retinoid X receptor/peroxisome proliferator-activated receptor- γ heterodimer. *J. Biol. Chem.* 282, 37738–37746.
- [28] Bai, P. et al. (2011) PARP-2 regulates SIRT1 expression and whole-body energy expenditure. *Cell Metab.* 13, 450–460.
- [29] Szanto, M. et al. (2011) Poly(ADP-ribose) polymerase-2 depletion reduces doxorubicin-induced damage through SIRT1 induction. *Cardiovasc. Res.* 92, 430–438.
- [30] Mehrotra, P., Riley, J.P., Patel, R., Li, F., Voss, L. and Goenka, S. (2011) PARP-14 functions as a transcriptional switch for Stat6-dependent gene activation. *J. Biol. Chem.* 286, 1767–1776.
- [31] Mehrotra, P., Hollenbeck, A., Riley, J.P., Li, F., Patel, R.J., Akhtar, N. and Goenka, S. (2012) Poly(ADP-ribose) polymerase 14 and its enzyme activity regulates T(H)2 differentiation and allergic airway disease. *J. Allergy Clin. Immunol.* 2012, 25. <<http://dx.doi.org/10.1016/j.jaci.2012.06.015>>.

- [32] Hassa, P.O., Covic, M., Hasan, S., Imhof, R. and Hottiger, M.O. (2001) The enzymatic and DNA binding activity of PARP-1 are not required for NF-kappa B coactivator function. *J. Biol. Chem.* 276, 45588–45597.
- [33] Hinz, M., Stilmann, M., Arslan, S.C., Khanna, K.K., Dittmar, G. and Scheidereit, C. (2010) A cytoplasmic ATM-TRAF6-clAP1 module links nuclear DNA damage signaling to ubiquitin-mediated NF-kappaB activation. *Mol. Cell* 40, 63–74.
- [34] Stilmann, M., Hinz, M., Arslan, S.C., Zimmer, A., Schreiber, V. and Scheidereit, C. (2009) A nuclear poly(ADP-ribose)-dependent signalosome confers DNA damage-induced IkkappaB kinase activation. *Mol. Cell* 36, 365–378.
- [35] Szanto, M., Brunyánszki, A., Kiss, B., Nagy, L., Gergely, P., Virag, L. and Bai, P. (2012) Poly(ADP-ribose) polymerase-2: emerging transcriptional roles of a DNA repair protein. *Cell. Mol. Life Sci.*, <http://dx.doi.org/10.1007/s00018-012-1003-8>.
- [36] Phulwani, N.K. and Kielian, T. (2008) Poly (ADP-ribose) polymerases (PARPs) 1–3 regulate astrocyte activation. *J. Neurochem.* 106, 578–590.
- [37] Popoff, I., Jijon, H., Monia, B., Tavernini, M., Ma, M., McKay, R. and Madsen, K. (2002) Antisense oligonucleotides to poly(ADP-ribose) polymerase-2 ameliorate colitis in interleukin-10-deficient mice. *J. Pharmacol. Exp. Ther.* 303, 1145–1154.
- [38] Zingarelli, B., Szabo, C. and Salzman, A.L. (1999) Blockade of poly(ADP-ribose) synthetase inhibits neutrophil recruitment, oxidant generation, and mucosal injury in murine colitis. *Gastroenterology* 116, 335–345.
- [39] Haddad, M., Rhinn, H., Bloquel, C., Coqueran, B., Szabo, C., Plotkine, M., Scherman, D. and Margail, I. (2006) Anti-inflammatory effects of PJ34, a poly(ADP-ribose) polymerase inhibitor, in transient focal cerebral ischemia in mice. *Br. J. Pharmacol.* 149, 23–30.
- [40] Espinoza, L.A., Smulson, M.E. and Chen, Z. (2007) Prolonged poly(ADP-ribose) polymerase-1 activity regulates JP-8-induced sustained cytokine expression in alveolar macrophages. *Free Radical Biol. Med.* 42, 1430–1440.
- [41] Lowenstein, C.J., Dinerman, J.L. and Snyder, S.H. (1994) Nitric oxide: a physiologic messenger. *Ann. Intern. Med.* 120, 227–237.
- [42] Moncada, S., Palmer, R.M. and Higgs, E.A. (1991) Nitric oxide: physiology, pathophysiology, and pharmacology. *Pharmacol. Rev.* 43, 109–142.
- [43] Ischiropoulos, H., Zhu, L. and Beckman, J.S. (1992) Peroxynitrite formation from macrophage-derived nitric oxide. *Arch. Biochem. Biophys.* 298, 446–451.
- [44] Ischiropoulos, H., Zhu, L., Chen, J., Tsai, M., Martin, J.C., Smith, C.D. and Beckman, J.S. (1992) Peroxynitrite-mediated tyrosine nitration catalyzed by superoxide dismutase. *Arch. Biochem. Biophys.* 298, 438–445.
- [45] Szabo, C., Ischiropoulos, H. and Radi, R. (2007) Peroxynitrite: biochemistry, pathophysiology and development of therapeutics. *Nat. Rev. Drug Discovery* 6, 662–680.
- [46] Xia, Y. and Zweier, J.L. (1997) Superoxide and peroxynitrite generation from inducible nitric oxide synthase in macrophages. *Proc. Natl. Acad. Sci. USA* 94, 6954–6958.
- [47] Mungrue, I.N. et al. (2002) Cardiomyocyte overexpression of iNOS in mice results in peroxynitrite generation, heart block, and sudden death. *J. Clin. Invest.* 109, 735–743.
- [48] Pacher, P., Beckman, J.S. and Liaudet, L. (2007) Nitric oxide and peroxynitrite in health and disease. *Physiol. Rev.* 87, 315–424.
- [49] Lin, Y., Tang, X., Zhu, Y., Shu, T. and Han, X. (2011) Identification of PARP-1 as one of the transcription factors binding to the repressor element in the promoter region of COX-2. *Arch. Biochem. Biophys.* 505, 123–129.
- [50] Le, N.T., Xue, M., Castelnoble, L.A. and Jackson, C.J. (2007) The dual personalities of matrix metalloproteinases in inflammation. *Front Biosci.* 12, 1475–1487.
- [51] Bai, P. et al. (2009) Poly(ADP-ribose) polymerase mediates inflammation in a mouse model of contact hypersensitivity. *J. Invest. Dermatol.* 129, 234–238.
- [52] Kauppinen, T.M. and Swanson, R.A. (2005) Poly(ADP-ribose) polymerase-1 promotes microglial activation, proliferation, and matrix metalloproteinase-9-mediated neuron death. *J. Immunol.* 174, 2288–2296.
- [53] Koh, S.H. et al. (2005) Effect of 3-aminobenzamide, PARP inhibitor, on matrix metalloproteinase-9 level in plasma and brain of ischemic stroke model. *Toxicology* 214, 131–139.
- [54] Oumouna-Benachour, K. et al. (2007) Poly(ADP-ribose) polymerase inhibition reduces atherosclerotic plaque size and promotes factors of plaque stability in apolipoprotein E-deficient mice: effects on macrophage recruitment, nuclear factor-kappaB nuclear translocation, and foam cell death. *Circulation* 115, 2442–2450.
- [55] Berger, N.A. (1985) Poly(ADP-ribose) in the cellular response to DNA damage. *Radiat. Res.* 101, 4–15.
- [56] Ying, W., Garnier, P. and Swanson, R.A. (2003) NAD⁺ repletion prevents PARP-1-induced glycolytic blockade and cell death in cultured mouse astrocytes. *Biochem. Biophys. Res. Commun.* 308, 809–813.
- [57] Ha, H.C. and Snyder, S.H. (1999) Poly(ADP-ribose) polymerase is a mediator of necrotic cell death by ATP depletion. *Proc. Natl. Acad. Sci. USA* 96, 13978–13982.
- [58] Tewari, M. et al. (1995) Yama/CPP32 beta, a mammalian homolog of CED-3, is a CrmA-inhibitable protease that cleaves the death substrate poly(ADP-ribose) polymerase. *Cell* 81, 801–809.
- [59] Benchoua, A. et al. (2002) Active caspase-8 translocates into the nucleus of apoptotic cells to inactivate poly(ADP-ribose) polymerase-2. *J. Biol. Chem.* 277, 34217–34222.
- [60] Cho, Y.S., Park, S.Y., Shin, H.S. and Chan, F.K. (2010) Physiological consequences of programmed necrosis, an alternative form of cell demise. *Mol. Cells* 29, 327–332.
- [61] Bai, P., Bakondi, E., Szabo, E., Gergely, P., Szabo, C. and Virag, L. (2001) Partial protection by poly(ADP-ribose) polymerase inhibitors from nitroxyl-induced cytotoxicity in thymocytes. *Free Radical Biol. Med.* 31, 1616–1623.
- [62] Virag, L., Salzman, A.L. and Szabo, C. (1998) Poly(ADP-ribose) synthetase activation mediates mitochondrial injury during oxidant-induced cell death. *J. Immunol.* 161, 3753–3759.
- [63] Bai, P., Hegedus, C., Erdelyi, K., Szabo, E., Bakondi, E., Gergely, S., Szabo, C. and Virag, L. (2007) Protein tyrosine nitration and poly(ADP-ribose) polymerase activation in N-methyl-N-nitro-N-nitrosoguanidine-treated thymocytes: implication for cytotoxicity. *Toxicol. Lett.* 170, 203–213.
- [64] Datta, R. et al. (2011) PARP-1 deficiency blocks IL-5 expression through calpain-dependent degradation of STAT-6 in a murine asthma model. *Allergy* 66, 853–861.
- [65] Kalogeropoulos, A.P., Georgiopoulou, V.V. and Butler, J. (2012) From risk factors to structural heart disease: the role of inflammation. *Heart Fail. Clin.* 8, 113–123.
- [66] Ungvari, Z., Gupte, S.A., Recchia, F.A., Batkai, S. and Pacher, P. (2005) Role of oxidative-nitrosative stress and downstream pathways in various forms of cardiomyopathy and heart failure. *Curr. Vasc. Pharmacol.* 3, 221–229.
- [67] Soriano, F. et al. (2001) Diabetic endothelial dysfunction: the role of poly(ADP-ribose) polymerase activation. *Nat. Med.* 7, 108–113.
- [68] Soriano, F.G., Pacher, P., Mabley, J., Liaudet, L. and Szabo, C. (2001) Rapid reversal of the diabetic endothelial dysfunction by pharmacological inhibition of poly(ADP-ribose) polymerase. *Circ. Res.* 89, 684–691.
- [69] Pacher, P., Liaudet, L., Mabley, J., Komjati, K. and Szabo, C. (2002) Pharmacologic inhibition of poly(adenosine diphosphate-ribose) polymerase may represent a novel therapeutic approach in chronic heart failure. *J. Am. Coll. Cardiol.* 40, 1006–1016.
- [70] Pacher, P., Liaudet, L., Mabley, J.G., Cziraki, A., Hasko, G. and Szabo, C. (2006) Beneficial effects of a novel ultrapotent poly(ADP-ribose) polymerase inhibitor in murine models of heart failure. *Int. J. Mol. Med.* 17, 369–375.
- [71] Pacher, P. and Szabo, C. (2008) Role of the peroxynitrite-poly(ADP-ribose) polymerase pathway in human disease. *Am. J. Pathol.* 173, 2–13. Epub 2008 Jun 5.
- [72] Virag, L., Szabo, E., Bakondi, E., Bai, P., Gergely, P., Hunyadi, J. and Szabo, C. (2002) Nitric oxide-peroxynitrite-poly(ADP-ribose) polymerase pathway in the skin. *Exp. Dermatol.* 11, 189–202.
- [73] Morrow, D.A. et al. (2009) A randomized, placebo-controlled trial to evaluate the tolerability, safety, pharmacokinetics, and pharmacodynamics of a potent inhibitor of poly(ADP-ribose) polymerase (INO-1001) in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention: results of the TIMI 37 trial. *J. Thromb. Thrombolysis* 27, 359–364.
- [74] Fong, P.C. et al. (2009) Inhibition of poly(ADP-Ribose) polymerase in tumors from BRCA mutation carriers. *N. Engl. J. Med.* 361 (2), 123–134.
- [75] Javle, M. and Curtin, N.J. (2012) The potential for poly (ADP-ribose) polymerase inhibitors in cancer therapy. *Ther. Adv. Med. Oncol.* 3, 257–267.
- [76] Hur, J.W., Sung, Y.K., Shin, H.D., Park, B.L., Cheong, H.S. and Bae, S.C. (2006) Poly(ADP-ribose) polymerase (PARP) polymorphisms associated with nephritis and arthritis in systemic lupus erythematosus. *Rheumatology (Oxford)* 45, 711–717.
- [77] Pascual, M., Lopez-Nevot, M.A., Caliz, R., Ferrer, M.A., Balsa, A., Pascual-Salcedo, D. and Martin, J. (2003) A poly(ADP-ribose) polymerase haplotype spanning the promoter region confers susceptibility to rheumatoid arthritis. *Arthritis Rheum.* 48, 638–641.
- [78] Mabley, J.G., Horvath, E.M., Murthy, K.G., Zsengeller, Z., Vaslin, A., Benko, R., Kollai, M. and Szabo, C. (2005) Gender differences in the endotoxin-induced inflammatory and vascular responses: potential role of poly(ADP-ribose) polymerase activation. *J. Pharmacol. Exp. Ther.* 315, 812–820.
- [79] Levaot, N. et al. (2011) Loss of Tankyrase-mediated destruction of 3BP2 is the underlying pathogenic mechanism of cherubism. *Cell* 147, 1324–1339.
- [80] Rapizzi, E., Fossati, S., Moroni, F. and Chiarugi, A. (2004) Inhibition of poly(ADP-ribose) glycohydrolase by galloctannin selectively up-regulates expression of proinflammatory genes. *Mol. Pharmacol.* 66, 890–898.
- [81] Genovese, T. et al. (2004) Treatment with a novel poly(ADP-ribose) glycohydrolase inhibitor reduces development of septic shock-like syndrome induced by zymosan in mice. *Crit. Care Med.* 32, 1365–1374.
- [82] Jagtap, P. et al. (2002) Novel phenanthridinone inhibitors of poly (adenosine 5'-diphosphate-ribose) synthetase: potent cytoprotective and antishock agents. *Crit. Care Med.* 30, 1071–1082.
- [83] Albertini, M., Clement, M.G., Lafortuna, C.L., Caniatti, M., Magder, S., Abdulmalek, K. and Hussain, S.N. (2000) Role of poly-(ADP-ribose) synthetase in lipopolysaccharide-induced vascular failure and acute lung injury in pigs. *J. Crit. Care* 15, 73–83.
- [84] Liaudet, L., Soriano, F.G., Szabo, E., Virag, L., Mabley, J.G., Salzman, A.L. and Szabo, C. (2000) Protection against hemorrhagic shock in mice genetically deficient in poly(ADP-ribose)polymerase. *Proc. Natl. Acad. Sci. USA* 97, 10203–10208.
- [85] Scott, G.S., Hake, P., Kean, R.B., Virag, L., Szabo, C. and Hooper, D.C. (2001) Role of poly(ADP-ribose) synthetase activation in the development of experimental allergic encephalomyelitis. *J. Neuroimmunol.* 117, 78–86.
- [86] Koedel, U., Winkler, F., Angele, B., Fontana, A. and Pfister, H.W. (2002) Meningitis-associated central nervous system complications are mediated by

- the activation of poly(ADP-ribose) polymerase. *J. Cereb. Blood Flow Metab.* 22, 39–49.
- [87] Kaplin, A.I. et al. (2005) IL-6 induces regionally selective spinal cord injury in patients with the neuroinflammatory disorder transverse myelitis. *J. Clin. Invest.* 115, 2731–2741.
- [88] Peralta-Leal, A., Rodriguez-Vargas, J.M., Aguilar-Quesada, R., Rodriguez, M.I., Linares, J.L., de Almodovar, M.R. and Oliver, F.J. (2009) PARP inhibitors: new partners in the therapy of cancer and inflammatory diseases. *Free Radical Biol. Med.* 47, 13–26.
- [89] Morisugi, T., Tanaka, Y., Kawakami, T. and Kirita, T. (2010) Mechanical stretch enhances NF-kappaB-dependent gene expression and poly(ADP-ribose) synthesis in synovial cells. *J. Biochem.* 147, 633–644.
- [90] Mota, R.A. et al. (2005) Inhibition of poly(ADP-ribose) polymerase attenuates the severity of acute pancreatitis and associated lung injury. *Lab. Invest.* 85, 1250–1262.
- [91] Soriano, F.G., Liaudet, L., Szabo, E., Virag, L., Mabley, J.G., Pacher, P. and Szabo, C. (2002) Resistance to acute septic peritonitis in poly(ADP-ribose) polymerase-1-deficient mice. *Shock* 17, 286–292.
- [92] Virag, L. et al. (2004) Effects of poly(ADP-ribose) polymerase inhibition on inflammatory cell migration in a murine model of asthma. *Med. Sci. Monit.* 10, BR77–BR83.
- [93] Boulares, A.H., Zoltoski, A.J., Sherif, Z.A., Jolly, P., Massaro, D. and Smulson, M.E. (2003) Gene knockout or pharmacological inhibition of poly(ADP-ribose) polymerase-1 prevents lung inflammation in a murine model of asthma. *Am. J. Respir. Cell Mol. Biol.* 28, 322–329.
- [94] Oumouna, M., Datta, R., Oumouna-Benachour, K., Suzuki, Y., Hans, C., Matthews, K., Fallon, K. and Boulares, H. (2006) Poly(ADP-ribose) polymerase-1 inhibition prevents eosinophil recruitment by modulating Th2 cytokines in a murine model of allergic airway inflammation: a potential specific effect on IL-5. *J. Immunol.* 177, 6489–6496.
- [95] Hageman, G.J., Larik, I., Pennings, H.J., Haenen, G.R., Wouters, E.F. and Bast, A. (2003) Systemic poly(ADP-ribose) polymerase-1 activation, chronic inflammation, and oxidative stress in COPD patients. *Free Radical Biol. Med.* 35, 140–148.
- [96] Cuzzocrea, S. et al. (2002) Pyrrolidine dithiocarbamate attenuates the development of acute and chronic inflammation. *Br. J. Pharmacol.* 135, 496–510.
- [97] Huang, H., McIntosh, J.L., Fang, L., Szabo, C. and Hoyt, D.G. (2003) Integrin-mediated suppression of endotoxin-induced DNA damage in lung endothelial cells is sensitive to poly(ADP-ribose) polymerase-1 gene deletion. *Int. J. Mol. Med.* 12, 533–540.
- [98] Kim, J.H. et al. (2008) Inflammatory and transcriptional roles of poly(ADP-ribose) polymerase in ventilator-induced lung injury. *Crit. Care* 12, R108.
- [99] Farivar, A.S. et al. (2004) Poly (ADP) ribose synthetase inhibition reduces obliterative airway disease in rat tracheal allografts. *J. Heart Lung Transplant.* 23, 993–1002.
- [100] Szabo, E. et al. (2001) Peroxynitrite production, DNA breakage, and poly(ADP-ribose) polymerase activation in a mouse model of oxazolone-induced contact hypersensitivity. *J. Invest. Dermatol.* 117, 74–80.
- [101] Farkas, B., Magyarlaki, M., Csete, B., Nemeth, J., Rablaczky, G., Bernath, S., Literati Nagy, P. and Sumegi, B. (2002) Reduction of acute photodamage in skin by topical application of a novel PARP inhibitor. *Biochem. Pharmacol.* 63, 921–932.
- [102] Mabley, J.G. et al. (2001) Anti-inflammatory effects of a novel, potent inhibitor of poly (ADP-ribose) polymerase. *Inflamm. Res.* 50, 561–569.
- [103] Shi, X. and Nuttall, A.L. (2007) Expression of adhesion molecular proteins in the cochlear lateral wall of normal and PARP-1 mutant mice. *Heart Res.* 224, 1–14.
- [104] Chiang, J. et al. (2009) Honokiol protects rats against eccentric exercise-induced skeletal muscle damage by inhibiting NF-kappaB induced oxidative stress and inflammation. *Eur. J. Pharmacol.* 610, 119–127.
- [105] Drel, V.R., Pacher, P., Stavniichuk, R., Xu, W., Zhang, J., Kuchmerovska, T.M., Slusher, B. and Obrosova, I.G. (2011) Poly(ADP-ribose)polymerase inhibition counteracts renal hypertrophy and multiple manifestations of peripheral neuropathy in diabetic Akita mice. *Int. J. Mol. Med.* 28, 629–635.
- [106] Mukhopadhyay, P., Horvath, B., Kechrid, M., Tanchian, G., Rajesh, M., Naura, A.S., Boulares, A.H. and Pacher, P. (2011) Poly(ADP-ribose) polymerase-1 is a key mediator of cisplatin-induced kidney inflammation and injury. *Free Radical Biol. Med.* 51, 1774–1788.
- [107] Shevalye, H. et al. (2010) Poly(ADP-ribose) polymerase (PARP) inhibition counteracts multiple manifestations of kidney disease in long-term streptozotocin-diabetic rat model. *Biochem. Pharmacol.* 79, 1007–1014.
- [108] Szabo, C., Biser, A., Benko, R., Bottinger, E. and Susztak, K. (2006) Poly(ADP-ribose) polymerase inhibitors ameliorate nephropathy of type 2 diabetic Leprd/db mice. *Diabetes* 55, 3004–3012.
- [109] Martinet, W., Knaapen, M.W., De Meyer, G.R., Herman, A.G. and Kockx, M.M. (2002) Elevated levels of oxidative DNA damage and DNA repair enzymes in human atherosclerotic plaques. *Circulation* 106, 927–932.
- [110] Benko, R., Pacher, P., Vaslin, A., Kollai, M. and Szabo, C. (2004) Restoration of the endothelial function in the aortic rings of apolipoprotein E deficient mice by pharmacological inhibition of the nuclear enzyme poly(ADP-ribose) polymerase. *Life Sci.* 75, 1255–1261.
- [111] Hans, C.P., Zerfaoui, M., Naura, A.S., Troxclair, D., Strong, J.P., Matrougui, K. and Boulares, A.H. (2009) Thieno[2,3-c]isoquinolin-5-one, a potent poly(ADP-ribose) polymerase inhibitor, promotes atherosclerotic plaque regression in high-fat diet-fed apolipoprotein E-deficient mice: effects on inflammatory markers and lipid content. *J. Pharmacol. Exp. Ther.* 329, 150–158.